Quantum propagator from Heisenberg equations

F. Kheirandish[1,*]

[1] University of Isfahan, Isfahan, Iran.

Quantum propagator for a general quadratic Lagrangian is obtained using position and momentum operators in Heisenberg picture and properties of propagators. The method can be generalized to a general quadratic Lagrangian in higher dimensions straightforwardly.

arXiv:1612.07674v1 [quant-ph] 22 Dec 2016

I. INTRODUCTION

The propagators are extensively used in many branches of Physics and interdisciplinary fields like quantum electrodynamics, quantum statistical mechanics, condensed matter Physics, polymer Physics, astrophysics and economics[1]. In non-relativistic quantum mechanics, propagators can be calculated in three different approaches[2]: (i) Using the explicit form of the energy eigenstates satisfying predefined boundary conditions. (ii) Solving the Heisenberg equations of motion for the problem in question. (iii) Using path integral techniques. The third approach from a pedagogical standpoint is interesting since it points out to the analogy with classical mechanics trough action principle. Due to the formal equivalence of the time-evolution operator in quantum mechanics and the density operator in quantum statistical mechanics, it turns out that the propagator formalism is also useful in calculating partition functions or thermodynamical quantities in quantum statistical mechanics.

In non-relativistic quantum mechanics the propagator gives the probability amplitude for a particle to travel from one spatial point at one time to another spatial point at a later time. This means that, if a system has Hamiltonian H, then the appropriate propagator is a function[3]

$$G(x,t;x',t') = \frac{1}{i\hbar}\Theta(t-t')K(x,t;x',t'), \qquad (1)$$

satisfying

$$(i\hbar\frac{\partial}{\partial t} - H_x)G(x,t;x',t') = \delta(x-x')\Theta(t-t'), \qquad (2)$$

where $\Theta(t)$ is the Heaviside step function and $K(x,t;x',t')$ is the kernel of the differential operator in question, often referred to as the propagator instead of G in this context. The propagator can also be written as

$$K(x,t;x',t') = \langle x|\hat{U}(t,t')|x'\rangle, \qquad (3)$$

where $\hat{U}(t,t')$ is the unitary time-evolution operator for the system taking states at time t to states at time t'.

In the present work we use the second approach to find the quantum propagator for a general quadratic Lagrangian in one spatial dimension. The generalization to higher dimensions is straightforward along the steps introduced here. This approach is based on the solution of the Heisenberg equation for the position and momentum operators and the properties of the propagator. To our knowledge this approach has not been discussed in standard text books on quantum mechanics and mathematically is a rigorous and simple method.

II. LAGRANGIAN

The general form of a classical quadratic Lagrangian is

$$L = \frac{1}{2}m\dot{x}^2 + a_1(t)x\dot{x} + \frac{1}{2}a_2(t)x^2 + a_3(t)\dot{x} + a_4(t)x. \qquad (4)$$

Regarding the fact that a total time derivative can be removed from Lagrangian, we rewrite the second and fourth terms as

$$\begin{aligned} a_1(t)x\dot{x} &= \frac{d}{dt}\big(\frac{1}{2}a_1(t)x^2\big) - \frac{1}{2}\dot{a}_1(t)x^2, \\ a_3(t)\dot{x} &= \frac{d}{dt}\big(a_3(t)x\big) - \dot{a}_3(t)x, \end{aligned} \qquad (5)$$

inserting Eq.(5) into Eq.(4) and defining new functions $c(t) = \dot{a}_1(t) - a_2(t)$ and $e(t) = \dot{a}_3(t) - a_4(t)$, we can rewrite Lagrangian Eq.(4) as[4]

$$L = \frac{1}{2}m\dot{x}^2 - \frac{1}{2}c(t)x^2 - e(t)x. \qquad (6)$$

The conjugate momentum is defined by

$$p = \frac{\partial L}{\partial \dot{x}} = m\dot{x}, \qquad (7)$$

and the corresponding Hamiltonian $H = p\dot{x} - L$, is

$$H = \frac{p^2}{2m} + \frac{1}{2}c(t)x^2 + e(t)x. \qquad (8)$$

The system is quantized by promoting classical variables (x,p) to quantum partners $(\hat{x},\hat{p})$ and imposing $[\hat{x},\hat{p}] = i\hbar$. From Heisenberg equations for position and momentum, we find

$$\ddot{\hat{x}} + \frac{c(t)}{m}\hat{x} = -\frac{e(t)}{m}. \qquad (9)$$

Equation (9) can be solved formally as

$$\hat{x}(t) = \hat{x}_H(t) - \frac{1}{m}\int_0^t dt'\, G(t,t')\, e(t'), \qquad (10)$$

where $G(t,t')$ is Green's function of Eq.(9) satisfying

$$\left[\frac{d^2}{dt^2}+\frac{1}{m}\,c(t)\right]G(t,t')=\delta(t-t'), \tag{11}$$

and $\hat{x}_H(t)$ is the homogeneous solution

$$\ddot{\hat{x}}_H(t)+\frac{c(t)}{m}\,\hat{x}_H(t)=0. \tag{12}$$

Let $f(t)$ and $g(t)$ be classical independent solutions of Eq.(12), then

$$\hat{x}_H(t)=\hat{A}f(t)+\hat{B}g(t), \tag{13}$$

where $\hat{A}$ and $\hat{B}$ are operator constants to be determined later. We can find the retarded Green's function in terms of the independent functions $f(t)$ and $g(t)$. The retarded Green's function is defined by

$$G(t,t')=\begin{cases}0, & t<t',\\ af(t)+bg(t), & t>t',\end{cases} \tag{14}$$

From the continuity and also the discontinuity of the time derivative of Green's function at $t=t'$, we will find

$$\begin{aligned} af(t')+bg(t')&=0,\\ \left.\frac{dG}{dt}\right|_{t=t'+}-\left.\frac{dG}{dt}\right|_{t=t'-}&=1, \end{aligned} \tag{15}$$

inserting Eq. (14) into Eq. (15), we find the retarded Green's function as

$$G(t,t')=\frac{g(t')f(t)-f(t')g(t)}{W(t')}\,\theta(t-t'), \tag{16}$$

where $\Theta(t-t')$ is Heaviside step function and $W(t)$ is Wronskian defined by

$$W(t)=\det\begin{pmatrix} g(t) & f(t)\\ \dot{g}(t) & \dot{f}(t)\end{pmatrix}. \tag{17}$$

From Eqs. (7,10,16), we have

$$\hat{x}(0) \;=\; \hat{x}_H(0)\equiv\hat{x}, \tag{18}$$

$$\hat{p}(0) \;=\; m\dot{x}(0)=m\dot{x}_H(0)\equiv\hat{p}. \tag{19}$$

The operator constants $\hat{A}$, $\hat{B}$ can be expressed in terms of the initial conditions or Schrödinger operators $\hat{x}$, $\hat{p}$ as

$$\begin{aligned} \hat{A} &= \frac{g(0)}{mW(0)}\,\hat{p}-\frac{\dot{g}(0)}{W(0)}\,\hat{x},\\ \hat{B} &= \frac{\dot{f}(0)}{W(0)}\,\hat{x}-\frac{f(0)}{mW(0)}\,\hat{p}. \end{aligned} \tag{20}$$

Now by inserting Eq.(20) into Eq.(13) and using Eq. (10), we find the the position operator as

$$\begin{aligned} \hat{x}(t) \;=\; & \frac{g(t)\dot{f}(0)-f(t)\dot{g}(0)}{W(0)}\,\hat{x}+\frac{f(t)g(0)-g(t)f(0)}{W(0)}\,\frac{\hat{p}}{m}\\ & -\frac{1}{m}\int_0^t dt'\,G(t,t')e(t'). \end{aligned} \tag{21}$$

III. PROPAGATOR

In Heisenberg picture, the time evolution of the position operator is given by

$$\hat{x}(t)=\hat{U}^\dagger(t)\hat{x}(0)\hat{U}(t), \tag{22}$$

or equivalently

$$\hat{U}(t)\hat{x}(t)=\hat{x}(0)\hat{U}(t). \tag{23}$$

Inserting Eq.(21) into Eq.(23) and taking the matrix elements in position space, we find

$$\langle x|U(t)[\alpha(t)x+\frac{\beta(t)}{m}p-\frac{\gamma(t)}{m}]|x'\rangle=x\langle x|U(t)|x'\rangle, \tag{24}$$

where

$$\alpha(t)=\frac{g(t)\dot{f}(0)-f(t)\dot{g}(0)}{W(0)}, \tag{25}$$

$$\beta(t)=\frac{f(t)g(0)-g(t)f(0)}{W(0)}, \tag{26}$$

$$\gamma(t)=\int_0^t dt'\,G(t,t')\,e(t'). \tag{27}$$

From Eq.(24) and the definition $K(x,t;x',0)=\langle x|U(t)|x'\rangle$, we find

$$\left(x'\alpha(t)+i\hbar\frac{\beta(t)}{m}\frac{\partial}{\partial x'}-\frac{\gamma(t)}{m}\right)K(x,t;x',0)=xK(x,t;x',0), \tag{28}$$

where we used $\hat{x}|x'\rangle=x'|x'\rangle$, $\hat{p}|x'\rangle=i\hbar\frac{\partial}{\partial x'}|x'\rangle$. Eq. (28) can be rewritten as

$$\frac{\partial}{\partial x'}\ln K(x,t;x',0)=-\frac{i}{\hbar}\frac{m}{\beta(t)}\,[x-x'\alpha(t)+\frac{\gamma(t)}{m}]. \tag{29}$$

Since the right side of Eq.(29) is linear in x', we can write

$$\ln K(x,t;x',0)=C_0+C_1x'+\frac{1}{2}C_2x'^2, \tag{30}$$

where C_0, C_1 and C_2 may depend on x and t. Inserting Eq.(30) into Eq.(29), we find

$$(C_1+C_2x')=-\frac{im}{\hbar\beta(t)}\left[x-\alpha(t)\,x'+\frac{\gamma}{m}\right], \tag{31}$$

and comparing the coefficients of x' on both sides we obtain

$$C_1=-\frac{im}{\hbar\beta}\,(x+\frac{\gamma}{m}),\quad C_2=\frac{im\alpha}{\hbar\beta}. \tag{32}$$

Therefore,

$$K(x,t;x',0)=e^{C_0(x,t)}e^{-\frac{im}{\hbar\beta}xx'}e^{\frac{im\alpha}{2\hbar\beta}x'^2}e^{-\frac{i\gamma}{\hbar\beta}x'}. \tag{33}$$

Now for the momentum operator we have

$$\hat{p}(t) = m\alpha(t)\hat{x} + \beta(t)\hat{p} - \gamma(t) = U^{\dagger}(t)\hat{p}U(t), \tag{34}$$

therefore, $\hat{U}(t)\hat{p}(t) = \hat{p}\hat{U}(t)$, leading to

$$\langle x|\hat{U}(t)[m\dot{\alpha}(t)\hat{x} + \dot{\beta}(t)\hat{p} - \dot{\gamma}(t)]|x'\rangle = \langle x|\hat{p}\hat{U}(t)|x'\rangle, \tag{35}$$

or equivalently

$$(m\dot{\alpha}x' + i\hbar\dot{\beta}\frac{\partial}{\partial x'} - \dot{\gamma})K(x,t;x',0) = -i\hbar\frac{\partial}{\partial x}K(x,t:x',0). \tag{36}$$

By inserting Eq.(33) into Eq. (36), we find

$$\frac{\partial C_0(x,t)}{\partial x} = \frac{i}{\hbar}[(m\dot{\alpha} + \frac{m}{\beta} - \frac{m\alpha\dot{\beta}}{\beta})x' + \frac{m\dot{\beta}}{\beta}x + \frac{\gamma\dot{\beta} - \dot{\gamma}\beta}{\beta}], \tag{37}$$

using the identity $\alpha\dot{\beta} - \dot{\alpha}\beta = 1$, (see definitions Eq. (37) can be solved as

$$C_0(x,t) = \frac{im}{2\hbar\beta}\left[\dot{\beta}x^2 + \frac{2}{m}(\gamma\dot{\beta} - \dot{\gamma}\beta)x\right] + \frac{i}{\hbar}\psi(t), \tag{38}$$

where $\psi(t)$ is an unknown function to be determined. The propagator now can be written as

$$K(x,t;x',0) = e^{\frac{i}{\hbar}\psi(t)}e^{\frac{im}{2\hbar\beta}\left[\dot{\beta}x^2 + \alpha x'^2 - 2xx' + \frac{2}{m}(\gamma\dot{\beta} - \dot{\gamma}\beta)x - \frac{2\gamma}{m}x'\right]}. \tag{39}$$

By making use of the identity

$$\langle x|\hat{U}(t)\hat{U}^{\dagger}(t)|x'\rangle = \delta(x - x'), \tag{40}$$

or equivalently

$$\int dx''\, \langle x|\hat{U}(t)|x''\rangle\langle x''|\hat{U}^{\dagger}(t)|x'\rangle = \delta(x - x'), \tag{41}$$

we find easily

$$e^{\frac{i}{\hbar}\psi(t)} = \sqrt{\frac{m}{2\pi\hbar\beta}}\, e^{\frac{i}{\hbar}\varphi(t)}, \tag{42}$$

where $\varphi(t)$ is a real function to be determined. Therefore, propagator can be written as

$$K(x,t;x',0) = \sqrt{\frac{m}{2\pi\hbar\beta}}\, e^{\frac{i}{\hbar}\varphi(t)}e^{\frac{im}{2\hbar\beta}\left[\alpha x'^2 + \dot{\beta}x^2 - 2(\alpha\dot{\beta} - \dot{\alpha}\beta)xx'\right]} \times e^{\frac{i}{\hbar\beta}\left[(\gamma\dot{\beta} - \dot{\gamma}\beta)x - \gamma x'\right]}. \tag{43}$$

Another well known property of the propagator is

$$\lim_{t\to 0} K(x,t;x',0) = \delta(x - x'), \tag{44}$$

from Eqs.(25,26,27) we have

$$\begin{aligned} \alpha(t) &\underset{t\to 0}{\to} 1, \\ \gamma(t) &\underset{t\to 0}{\to} 0, \\ \beta(t) &\underset{t\to 0}{\to} t + o(t^2), \end{aligned} \tag{45}$$

where $\beta(t)$ given in Eq.(26) has been Taylor expanded around $t = 0$ up to the first order. Note that $\alpha(t)$ and $\gamma(t)$ in Eq.(43) can be simply replace by their respective limiting values but $\beta(t)$ carrying the singularity, should be replaced with its asymptotic value, namely t. Therefore,

$$\begin{aligned} \lim_{t\to 0} K(x,t;x',0) &= \lim_{t\to 0}\frac{e^{\frac{i}{\hbar}\varphi(0)}}{\sqrt{2\pi\hbar t}}\, e^{\frac{i}{2\hbar t}(x - x')^2}, \\ &= \delta(x - x'). \end{aligned} \tag{46}$$

Comparing this result with the identity

$$\lim_{t\to 0}(A/\pi t)^{\frac{1}{2}}e^{-\frac{A}{t}(x - x')^2} = \delta(x - x'), \tag{47}$$

we will find $e^{\frac{i}{\hbar}\varphi(0)} = 1/\sqrt{i}$. If we redefine $\lambda(t) = \varphi(t) - \varphi(0)$, then for the quantum propagator we find

$$K(x,t;x',0) = \sqrt{\frac{m}{2\pi i\hbar\beta}}\, e^{\frac{i}{\hbar}\lambda(t)}e^{\frac{im}{2\hbar\beta}\left[\dot{\beta}x^2 + \alpha x'^2 - 2xx' + \frac{2}{m}(\gamma\dot{\beta} - \dot{\gamma}\beta)x - \frac{2\gamma}{m}x'\right]}. \tag{48}$$

To determine $\lambda(t)$ we know that the propagator $K(x,t;0,0)$ satisfies Schrödinger equation

$$i\hbar\partial_t K(x,t;0,0) = \left[-\frac{\hbar^2}{2m}\partial_x^2 + \frac{1}{2}c(t) + e(t)x\right]K(x,t;0,0), \tag{49}$$

taking temporal and spatial differentiations, and setting $x = 0$, we find

$$\dot{\lambda}(t) = -\frac{1}{2m}\left(\frac{\gamma\dot{\beta} - \dot{\gamma}\beta}{\beta}\right)^2, \tag{50}$$

Finally, the quantum propagator corresponding to the quadratic Lagrangian Eq.(6) is

$$K(x,t;x',0) = \sqrt{\frac{m}{2\pi i\hbar\beta}}\, e^{-\frac{i}{2m\hbar}\int_0^t dt'\,(\frac{\gamma\dot{\beta} - \dot{\gamma}\beta}{\beta})^2} \times e^{\frac{im}{2\hbar\beta}\left[\dot{\beta}x^2 + \alpha x'^2 - 2xx' + \frac{2}{m}(\gamma\dot{\beta} - \dot{\gamma}\beta)x - \frac{2\gamma}{m}x'\right]}. \tag{51}$$

IV. A DIFFERENTIAL EQUATION FOR $\beta(t)$

Setting $t' = 0$ in Eq.(16) and comparing the result with Eq.(26), we have

$$G(t) = m\beta(t)\,\Theta(t). \tag{52}$$

From Eq.(26) it is clear that $\beta(t)$ satisfies the equation

$$\left[\frac{d^2}{dt^2} + \frac{1}{m}\,c(t)\right]\beta(t) = 0, \tag{53}$$

with the initial conditions

$$\beta(0) = 0, \quad \left.\frac{d\beta(t)}{dt}\right|_{t=0} = 1, \tag{54}$$

which is the same result obtained by Schulman[4] using a different approach.

Example 1. For a harmonic oscillator in an external classical source, we set $c(t) = m\omega^2$ and $e(t) \neq 0$, and find

$$\begin{aligned}
f(t) &= \cos(\omega t),\\
g(t) &= \sin(\omega t),\\
\alpha(t) &= \cos(\omega t),\\
\beta(t) &= \frac{\sin(\omega t)}{\omega},\\
G(t,t') &= \frac{\sin[\omega(t-t')]}{\omega}\,\theta(t-t'),\\
\gamma(t) &= \int_0^t dt'\,\frac{\sin[\omega(t-t')]}{\omega}\,e(t'),
\end{aligned}$$

therefore,

$$\begin{aligned}
&K(x,t;x',0) =\\
&\sqrt{\frac{m\omega}{2\pi i\hbar\sin(\omega t)}}\,e^{\frac{im\omega}{2\hbar\sin(\omega t)}\left[(x^2+x'^2)\cos(\omega t)-2xx'\right]}\\
&\times e^{-\frac{i}{\hbar\sin(\omega t)}\left[x\int_0^t dt'\,\sin(\omega t')e(t')+x'\int_0^t dt'\,\sin\omega(t-t')e(t')\right]}\\
&\times e^{-\frac{i}{m\hbar\omega\sin(\omega t)}\int_0^t dt'\int_0^{t'} ds\,e(t')e(s)\sin\omega(t-t')\sin(\omega s)}.
\end{aligned} \tag{55}$$

as expected[4].

V. CONCLUSION

Using position and momentum operators in Heisenberg picture and properties of propagators, quantum propagator for a general quadratic Lagrangian obtained. The method is mathematically rigorous and can be generalised to a general quadratic Lagrangian in higher dimensions straightforwardly.

VI. SUGGESTED PROBLEM

Consider the following quadratic Lagrangian in n spatial dimensions

$$L = \frac{1}{2}M\sum_{j=1}^{n}\dot{X}_j^2 - \frac{1}{2}\sum_{i,j=1}^{n}X_iB_{ij}X_j + \sum_{j=1}^{n}C_j(t)X_j, \tag{56}$$

where $B_{ij} = B_{ji}$ is a time independent symmetric matrix and $C_j(t)$ is a time dependent n-dimensional vector. Find the corresponding quantum propagator.

* Electronic address: fkheirandish@yahoo.com

[1] G. Barton, *Elements of Green's Functions and Propagation: Potentials, Diffusion, and Waves* (Oxford University Press, 1989).

[2] L. A. Beauregard, Am. J. Phys. 34, 324 (1966).

[3] J. J. Sakurai, *Modern Quantum Mechanics* (Addison-Wesley, Reading, Mass. 1994).

[4] L. S. Schulman, *Techniques and Applications of Path Integration*, (Dover Publications, Inc., 31 East 2nd Street, Mineola, N.Y. 11501).